

Name : _____

Grade : VIII

Subject : Mathematics

Chapter : 9. Algebraic Expressions and Identities

Objective Type Questions

(1 Marks)

I. Multiple choice questions

1. The product of a monomial and a binomial is a (NCERT Exemplar)
 a. monomial b. binomial c. trinomial d. none of these
2. If we subtract $-3x^2y^2$ from x^2y^2 , then we get (NCERT Exemplar)
 a. $-4x^2y^2$ b. $-2x^2y^2$ c. $2x^2y^2$ d. $4x^2y^2$
3. Product of the following monomials $4p$, $-7q^3$, $-7pq$ is (NCERT Exemplar)
 a. $196 p^2 q^4$ b. $196 pq^4$ c. $-196 p^2 q^4$ d. $196 p^2 q^3$
4. Area of a rectangle with length $4ab$ and breadth $6b^2$ is (NCERT Exemplar)
 a. $24 a^2 b^2$ b. $24 ab^3$ c. $24 ab^2$ d. $24 ab$
5. Volume of a rectangular box (cuboid) with length = $2ab$, breadth = $3ac$ and height = $2ac$ is (NCERT Exemplar)
 a. $12 a^3 bc^2$ b. $12 a^3 bc$ c. $12 a^2 bc$ d. $2ab + 3ac + 2ac$
6. What is the volume of the cuboid of length $8xy$, breadth $3xy$ and height xy ? (NCERT Exemplar)
 a. $5x^3 y^2$ b. $6x^3 y^3$ c. $24 x^3 y^3$ d. $8 x^2 y^3$
7. What is the value of $(x - y)^2$? (NCERT Exemplar)
 a. $x^2 + 2xy + y^2$ b. $x^2 - 2xy + y^2$ c. $x^2 - 2xy - y^2$ d. $x^2 + y^2$
8. The value of $(-27x^2y) \div (-9xy)$ is (NCERT Exemplar)
 a. $3xy$ b. $-3xy$ c. $-3x$ d. $3x$
9. The value of $(3x^3 + 9x^2 + 27x) \div 3x$ is (NCERT Exemplar)
 a. $x^2 + 9 + 27x$ b. $3x^3 + 3x^2 + 27x$ c. $3x^3 + 9x^2 + 9$ d. $x^3 + 3x + 9$
10. Which of the following is an identity? (NCERT Exemplar)
 a. $(p + q)^2 = p^2 + q^2$ b. $p^2 - q^2 = (p - q)^2$
 c. $p^2 - q^2 = p^2 + 2pq - q^2$ d. $(p + q)^2 = p^2 + 2pq + q^2$

11. The product of $-a$, $-a^2$ and a^3 equals to:

a. $-a^6$

b. a^6

c. $a - a^2 + a^3$

d. $a + a^2 + a^3$

(NCERT Exemplar)

12. Dividing $x^3 + 2x^2 + x$ by $x(x + 1)$, we get:

a. x

b. $x + 1$

c. $x + 2$

d. $x(x + 1)$

(NCERT Exemplar)

13. $a^2 - b^2$ is equal to:

a. $(a - b)^2$

b. $(a - b)(a - b)$

c. $(a + b)(a - b)$

d. $(a + b)(a + b)$

(NCERT Exemplar)

14. Which of the following are like terms?

a. $5xyz^2, -3xy^2z$

b. $-5xyz^2, 7xyz^2$

c. $5xyz^2, 5x^2yz$

d. $5xyz^2, x^2y^2z^2$

(NCERT Exemplar)

15. The value of $(a + b)^2 + (a - b)^2$ is

a. $2a + 2b$

b. $2a - 2b$

c. $2a^2 + 2b^2$

d. $2a^2 - 2b^2$

(NCERT Exemplar)

16. Product of $6a^2 - 7b + 5ab$ and $2ab$ is

a. $12a^3b - 14ab^2 + 10ab$

b. $12a^3b - 14ab^2 + 10a^2b^2$

c. $6a^2 - 7b + 7ab$

d. $12a^2b - 7ab^2 + 10ab$

(NCERT Exemplar)

17. Square of $3x - 4y$ is

a. $9x^2 - 16y^2$

b. $6x^2 - 8y^2$

c. $9x^2 + 16y^2 + 24xy$

d. $9x^2 + 16y^2 - 24xy$

(NCERT Exemplar)

18. Square of $9x - 7xy$ is

a. $81x^2 + 49x^2y^2$

b. $81x^2 - 49x^2y^2$

c. $81x^2 + 49x^2y^2 - 126x^2y$

d. $81x^2 + 49x^2y^2 - 63x^2y$

(NCERT Exemplar)

19. $a(b + c) = ab + bc$ is

a. Commutative property

b. Distributive property

c. Associative property

d. Closure property

1. b	2. d	3. a	4. b	5. a	6. c	7. b	8. d	9. d	10. d
11. b	12. b	13. c	14. b	15. c	16. b	17. d	18. c	19. b	

III. Multiple choice questions

1. Which is the like term as $24a^2bc$

(NCERT Exemplar)

a. $13 \times 8a \times 2b \times c \times a$

b. $8 \times 3 \times a \times b \times c$

c. $3 \times 8 \times a \times b \times c \times c$

d. $3 \times 8 \times a \times a \times b \times c$

2. Which of the following is correct?

(NCERT Exemplar)

a. $(a - b)^2 = a^2 + 2ab - b^2$

b. $(a - b)^2 = a^2 - 2ab + b^2$

c. $(a - b)^2 = a^2 - b^2$

d. $(a + b)^2 = a^2 + 2ab - b^2$

3. Which of the following is a binomial?

(NCERT Exemplar)

- a. $7 \times a + a$ b. $6a^2 + 7b + 2c$ c. $4a \times 3b \times 2c$ d. $6(a^2 + b)$

4. Coefficient of y in the term $\frac{-y}{3}$ is

(NCERT Exemplar)

- a. -1 b. -3 c. $\frac{-1}{3}$ d. $\frac{1}{3}$

5. The sum of $-7 pq$ and $2pq$ is

(NCERT Exemplar)

- a. $-9pq$ b. $9pq$ c. $5pq$ d. $-5pq$

6. The product of a monomial and a binomial is a

(NCERT Exemplar)

- a. monomial b. binomial c. trinomial

d. None of these

7. Sum of $a - b + ab$, $b + c - bc$ and $c - a - ac$ is

(NCERT Exemplar)

- a. $2c + ab - ac - bc$ b. $2c - ab - ac - bc$ c. $2c + ab + ac + bc$ d. $2c - ab + ac + bc$

1. a	2. b	3. d	4. c	5. d	6. b	7. a
------	------	------	------	------	------	------

I. Fill in the blanks

1. $38x^3y^2z \div 19xy^2$ is equal is _____.

(NCERT Exemplar)

2. Volume of a rectangular box with $l = b = h = 2x$ is _____.

(NCERT Exemplar)

3. The coefficient in $-37 abc$ is _____.

(NCERT Exemplar)

4. The sum of areas of two squares with sides $4a$ and $4b$ is _____.

(NCERT Exemplar)

5. The factorisation of $2x + 4y$ is _____.

(NCERT Exemplar)

1. $2x^2z$	2. $8x^3$	3. -37	4. $16(a^2 + b^2)$	5. $2(x + 2y)$
------------	-----------	--------	--------------------	----------------

I. True or False

1. $(a + b)(a - b) = a^2 - b^2$

(NCERT Exemplar)

2. The product of two negative term is a negative term.

(NCERT Exemplar)

3. The coefficient of the term $-6x^2y^2$ is -6.

(NCERT Exemplar)

4. The value of p for $51^2 - 49^2 = 100p$ is 2.

(NCERT Exemplar)

5. The value of $(a + 1)(a - 1)(a^2 + 1)$ is $a^4 - 1$.

(NCERT Exemplar)

1. True	2. False	3. True	4. True	5. True
---------	----------	---------	---------	---------

I. Very Short Answer Questions

1. Add the following.

[NCERT Exemplar]

a. $7a^2bc, -3abc^2, 3a^2bc, 2abc^2$

b. $5x^2 - 3xy + 4y^2 - 9, 7y^2 + 5xy - 2x^2 + 13$

Sol. a. $7a^2bc + (-3abc^2) + 3a^2bc + 2abc^2$

$$\Rightarrow 7a^2bc - 3abc^2 + 3a^2bc + 2abc^2$$

$$\Rightarrow 10a^2bc - abc^2$$

b. $[5x^2 - 3xy + 4y^2 - 9] + [7y^2 + 5xy - 2x^2 + 13]$

$$\Rightarrow 5x^2 - 3xy + 4y^2 - 9 + 7y^2 + 5xy - 2x^2 + 13$$

$$\Rightarrow 3x^2 + 11y^2 + 2xy + 4$$

2. Subtract $6x^2 - 4xy + 5y^2$ from $8y^2 + 6xy - 3x^2$

[NCERT Exemplar]

Sol.
$$\begin{array}{r} 8y^2 + 6xy - 3x^2 \\ 5y^2 - 4xy + 6x^2 \\ \hline - \quad + \quad - \\ 3y^2 + 10xy - 9x^2 \end{array}$$

3. Multiply the following.

a. $15xy^2, 17yz^2$

[NCERT Exemplar]

b. $-5a^2bc, 11ab, 13abc^2$

Sol. a. $15xy^2, 17yz^2 = (15 \times 17) \times x \times y^2 \times y \times z^2$

$$= 255 xy^3 z^2$$

b. $-5a^2bc \times 11ab \times 13abc^2 = (-5 \times 11 \times 13) a^2bc \times ab \times abc^2$

$$= -715 a^4 b^3 c^3$$

4. Find the product of $-4p$ and $7pq$.

Sol. We have,

$$-4p \times 7pq = (-4 \times 7) \times p \times pq$$

$$= -28p^2q$$

5. Simplify: $\left(\frac{3}{4}x - \frac{4}{3}y\right)^2 + 2xy$

[NCERT Exemplar]

Sol. $\left(\frac{3}{4}x - \frac{4}{3}y\right)^2 + 2xy$

$$\Rightarrow \left(\frac{3}{4}x\right)^2 - 2\left(\frac{3}{4}x\right)\left(\frac{4}{3}y\right) + \left(\frac{4}{3}y\right)^2 + 2xy$$

$$\Rightarrow \frac{9}{16}x^2 - 2xy + \frac{16}{9}y^2 + 2xy$$

$$\Rightarrow \frac{9}{16}x^2 + \frac{16}{9}y^2$$

6. Using suitable identify evaluate: $(35.4)^2 - (14.6)^2$.

[NCERT Exemplar]

Sol. $(35.4)^2 - (14.6)^2 = (35.4 + 14.6)(35.4 - 14.6)$

$$= 50 \times 20.8$$

$$= 1040$$

7. Expand the following using suitable identities.

a. $\left(\frac{2}{3}x - \frac{3}{2}y\right)^2$

b. $(7x + 5)^2$

Sol. $\left(\frac{2}{3}x - \frac{3}{2}y\right)^2$

$$\Rightarrow \left(\frac{2}{3}x\right)^2 - 2\left(\frac{2}{3}x\right)\left(\frac{3}{2}y\right) + \left(\frac{3}{2}y\right)^2$$

$$\Rightarrow \frac{4}{9}x^2 - 2xy + \frac{9}{4}y^2$$

b. $(7x + 5)^2$

$$\Rightarrow (7x)^2 + 2(7x)(5) + (5)^2$$

$$\Rightarrow 49x^2 + 70x + 25$$

8. Simplify : $(ab - c)^2 + 2abc$

[NCERT Exemplar]

Sol. $(ab - c)^2 + 2abc$

$$\Rightarrow (ab)^2 - 2(ab)(c) + (c)^2 + 2abc$$

$$\Rightarrow a^2b^2 - 2abc + c^2 + 2abc$$

$$\Rightarrow a^2b^2 + c^2$$

9. Find the product of $(-3x^2y) \times (4x^2y - 3xy^2 + 4x - 5y)$

Sol. $(-3x^2y) \times (4x^2y - 3xy^2 + 4x - 5y)$

$$= -12x^4y^2 + 9x^3y^3 - 12x^3y + 15x^2y^2$$

10. If the sides of a triangle are $3x + 1$, $-x + 2$ and $4x + 6$, then find its perimeter.

Sol. Let

AB = $3x + 1$

BC = $-x + 2$

AC = $4x + 6$

Since, perimeter of $\triangle ABC = AB + BC + AC$

$$= Bx + 1 + (-x + 2) + 4x + 6$$

$$= 3x + 1 - x + 2 + 4x + 6$$

$$= 6x + 9$$

II. Very Short Answer Questions

1. Using suitable identity, evaluate $(69.3)^2 - (30.7)^2$

(NCERT Exemplar)

Sol. $(69.3)^2 - (30.7)^2$

Using identify $a^2 - b^2 = (a + b)(a - b)$

$$(69.3)^2 - (30.7)^2 = (69.3 + 30.7)(69.3 - 30.7)$$

$$= 100 \times 38.6$$

$$= 3860$$

2. State whether the statements are True (T) or False (F).

(NCERT Exemplar)

i. The product of two negative term is a negative term.

ii. $p^2q + q^2r + r^2q$ is a binomial.

Sol. i. False

ii. False

3. Find the volume of a rectangular box with $l = b = h = 2x$

Sol. Volume of rectangular box = $l \times b \times h$

$$= 2x \times 2x \times 2x = 2 \times 2 \times 2 \times x \times x \times x = 8x^3$$

4. Add : $9ax, 3by - cz$ and $-5by + ax + 3cz$

(NCERT Exemplar)

Sol. $9ax$

$$3by - cz$$

$$+ ax - 5by + 3cz$$

$$\underline{10ax - 2by + 2cz}$$

6. Fill in the blanks.

i. Area of rectangular plot with sides $4x^2$ and $3y^2$ is _____.

ii. $a^2 - b^2 = (a + b) \underline{\hspace{2cm}}$.

Sol. i. $4x^2 \times 3y^2 = 12x^2y^2$

ii. $(a - b)$

7. Is 1 a polynomial? If yes, what is its degree?

Sol. Yes, 1 is a polynomial. Its degree is zero. (As it contains no variable)

I. Short Answer Questions

1. If $x - \frac{1}{x} = 7$, then find the value $x^2 + \frac{1}{x^2}$.

[NCERT Exemplar]

Sol. We know that $(a - b)^2 = a^2 - 2ab + b^2$

$$\therefore \left(x - \frac{1}{x}\right)^2 = 7^2$$

$$x^2 + \frac{1}{x^2} - 2 \times x \times \frac{1}{x} = 49$$

$$x^2 + \frac{1}{x^2} - 2 = 49$$

$$x^2 + \frac{1}{x^2} = 49 + 2$$

$$x^2 + \frac{1}{x^2} = 51$$

2. Simplify

a. $-pqr(p^2 + q^2 + r^2)$

b. $(px + qy)(ax - by)$

Sol. a. $-pqr(p^2 + q^2 + r^2)$

$$\begin{aligned} &= -(pqr) \times p^2 - (pqr) \times q^2 - (pqr) \times r^2 \\ &= -p^3 qr - pq^3 r - pqr^3 \end{aligned}$$

b. $(px + qy)(ax - by)$

$$\begin{aligned} &= px(ax - by) + qy(ax - by) \\ &= apx^2 - pbxy + aqxy - qby^2 \end{aligned}$$

3. Simplify:

a. $(3x + 2y)^2 - (3x - 2y)^2$

$$\begin{aligned} &\Rightarrow [(3x)^2 + 2(3x)(2y) + (2y)^2] - [(3x)^2 - 2(3x)(2y) + (2y)^2] \\ &\Rightarrow (9x^2 + 12xy + 4y^2) - (9x^2 - 12xy + 4y^2) \end{aligned}$$

$$\Rightarrow 9x^2 + 12xy + 4y^2 - 9x^2 + 12xy - 4y^2$$

$$\Rightarrow 24xy$$

$$\begin{aligned}
 & b. \left(\frac{7}{9}a + \frac{9}{7}b\right)^2 - ab \\
 & \Rightarrow \left(\frac{7}{9}a\right)^2 + 2\left(\frac{7}{9}a\right)\left(\frac{9}{7}b\right) + \left(\frac{9}{7}b\right)^2 - ab \\
 & \Rightarrow \frac{49}{81}a^2 + 2ab + \frac{81}{49}b^2 - ab \\
 & \Rightarrow \frac{49}{81}a^2 + ab + \frac{81}{49}b^2
 \end{aligned}$$

4. Simplify: $(2x + 5)^2 - (2x - 5)^2$

Sol. $(2x + 5)^2 - (2x - 5)^2 = [2x]^2 + (5)^2 + 2 \times 2x \times 5]$
 $- [(2x)^2 + (5)^2 - 2 \times 2x \times 5]$
 $= [4x^2 + 25 + 20x] - [4x^2 + 25 - 20x]$
 $= 4x^2 + 25 + 20x - 4x^2 + 25 - 20x$
 $= 20x + 20x$
 $= 40x$

5. Find the value of $\frac{38^2 - 22^2}{16}$, using a suitable identify.

[NCERT Exemplar]

Sol. Since, $a^2 - b^2 = (a + b)(a - b)$, therefore

$$\begin{aligned}
 38^2 - 22^2 &= (38 - 22)(38 + 22) \\
 &= 16 \times 60
 \end{aligned}$$

$$\begin{aligned}
 \text{So, } \frac{38^2 - 22^2}{16} &= \frac{16 \times 60}{16} \\
 &= 60
 \end{aligned}$$

6. Find the value of x , if $10000x = (9982)^2 - (18)^2$

[NCERT Exemplar]

Sol. R.H.S = $(9982)^2 - (18)^2$
 $= (9982 + 18)(9982 - 18)$
 $[$ Since, $a^2 - b^2 = (a + b)(a - b)$ $]$
 $= (10000) \times (9964)$

$$\text{L.H.S} = (10000) \times x$$

Comparing L.H.S. and R.H.S, we get

$$10000x = 10000 \times 9964$$

or

$$x = \frac{10000 \times 9964}{10000} = 9964$$

7. Verify that:

$$(3x + 5y)^2 - 30xy = 9x^2 + 25y^2$$

[NCERT Exemplar]

Sol. L.H.S. = $(3x + 5y)^2 - 30xy$

$$= (3x)^2 + 2 \times 3x \times 5y + (5y)^2 - 30xy$$

[Since, $(a + b)^2 = a^2 + 2ab + b^2$]

$$= 9x^2 + 30xy + 25y^2 - 30xy$$

$$= 9x^2 + 25y^2$$


$$= R.H.S$$

(Hence Verified)

8. Two adjacent sides of a rectangle are $5x^2 - 3y^2$ and $x^2 + 2xy$. Find the perimeter.

Sol. Let, Length, $l = x^2 + 2xy$

and Breadth, $b = 5x^2 - 3y^2$



Since, perimeter of rectangle = $2(l + b)$

Then, perimeter = $2[x^2 + 2xy + 5x^2 - 3y^2]$

$$= 2[6x^2 + 2xy - 3y^2]$$


$$= 12x^2 + 4xy - 6y^2$$

9. Multiply: $(x^4 + \frac{1}{x^4})$ by $(x + \frac{1}{4})$.

Sol. $(x^4 + \frac{1}{x^4})$ by $(x + \frac{1}{4}) = x^2 \times x + x^4 + \frac{1}{x} + \frac{1}{x^4} \times x + \frac{1}{x^4} \times \frac{1}{x}$

$$= x^5 + x^3 + \frac{1}{x^3} + \frac{1}{x^5}$$

10. Find the length of the side of the given square, if area of the square is 625 square units and then find the value of x .



[NCERT Exemplar]

Sol. We know that,

$$\text{Area of square} = (\text{side})^2$$

$$\begin{aligned} \Rightarrow & 625 = (4x + 5)^2 \\ \Rightarrow & 25 = 4x + 5 \\ \Rightarrow & 4x = 25 - 5 \\ \Rightarrow & 4x = 20 \\ \Rightarrow & x = \frac{20}{4} = 5 \end{aligned}$$

II. Short Answer Questions

1. Subtract $(x^3 - 5y^3 + 2z^3 + 3xyz)$ from the sum of $(-3x^3 - 7y^3 + z^3 + xyz)$:

$(3x^3 - y^3 + 7z^3 + 5xyz)$ and $(-x^3 - y^3 - 2x^3 - 3xyz)$.

$$\begin{aligned} \text{Sol. Let } A &= (-3x^3 - 7y^3 + z^3 + xyz) + (3x^3 - y^3 + 7z^3 + 5xyz) + (-x^3 - y^3 - 2z^3 - 3xyz) \\ &= (-3x^3 + 3x^3 - x^3) + (-7y^3 - y^3 - y^3) + (z^3 + 7z^3 - 2z^3) + (xyz + 5xyz - 3xyz) \\ &= -x^3 - 9y^3 + 6z^3 + 3xyz \end{aligned}$$

$$\text{Let } B = x^3 - 5y^3 + 2z^3 + 3xyz$$

For $A - B$, $-x^3 - 9y^3 + 6z^3 + 3xyz$ [By column method]

$$+ x^3 - 5y^3 + 2z^3 + 3xyz$$

$$\begin{array}{r} - \quad + \quad - \quad - \\ \hline -2x^3 - 4y^3 + 4z^3 + 0 \end{array}$$

$$\therefore A - B = -2x^3 - 4y^3 + 4z^3$$

2. Simplify the following expression.

$$x + [2x^2 - \{x^2 - 2xy - (-5x^2 - x)\}]$$

Sol. We have,

$$\begin{aligned} & x + [2x^2 - \{x^2 - 2xy - (-5x^2 - x)\}] \\ &= x + [2x^2 - \{x^2 - 2xy + 5x^2 + x\}] \\ &= x + [2x^2 - \{6x^2 + 2xy + x\}] \\ &= x + [2x^2 - 6x^2 + 2xy - x] \\ &= x + [-4x^2 + 2xy - x] \\ &= x - 4x^2 + 2xy - x = -4x^2 + 2xy \end{aligned}$$

Note: For simplification of algebraic expression, we follow same rules as in arithmetic and we also follow BDMAS / BODMAS and other brackets rules.

3. Subtract : $\frac{8}{5}x^3 - \frac{2}{3}x^2 + \frac{3}{2}xy - 2y^3$ from $\frac{1}{5}x^3 + \frac{3}{5}xy - \frac{1}{3}x^2 - 5y^3$

Sol. We have,

$$\begin{aligned} & \left(\frac{1}{5}x^3 + \frac{3}{5}xy - \frac{1}{3}x^2 - 5y^3 \right) - \left(\frac{8}{5}x^3 - \frac{2}{3}x^2 + \frac{3}{2}xy - 2y^3 \right) \\ &= \left(\frac{1}{5}x^3 + \frac{3}{5}xy - \frac{1}{3}x^2 - 5y^3 \right) + \left(-\frac{8}{5}x^3 + \frac{2}{3}x^2 - \frac{3}{2}xy + 2y^3 \right) \\ &= \left(\frac{1}{5}x^3 - \frac{8}{5}x^3 \right) + \left(-\frac{1}{3}x^2 + \frac{2}{3}x^2 \right) + \left(\frac{3}{5}xy - \frac{3}{2}xy \right) + (-5y^3 + 2y^3) \\ &= \left(\frac{1}{5} - \frac{8}{5} \right)x^3 + \left(\frac{-1+2}{3} \right)x^2 + \left(\frac{3}{5} - \frac{3}{2} \right)xy + (-5 + 2)y^3 \\ &= \left(\frac{1-8}{5} \right)x^3 + \left(\frac{-1+2}{3} \right)x^2 + \left(\frac{6-15}{10} \right)xy - 3y^3 \\ &= \frac{-7}{5}x^3 + \left(\frac{1}{3} \right)x^2 + \left(\frac{-9}{10} \right)xy - 3y^3 \\ &= -\frac{7}{5}x^3 + \frac{1}{3}x^2 - \frac{9}{10}xy - 3y^3 \end{aligned}$$

4. Expand the following, using suitable identities.

[NCERT Exemplar]

i. $\left(\frac{4x}{5} + \frac{y}{4} \right) \left(\frac{4x}{5} + \frac{3y}{4} \right)$

ii. $(x^2 + y^2)(x^2 - y^2)$

iii. $(0.9p - 0.5q)^2$

Sol. i. $\left(\frac{4x}{5} + \frac{y}{4} \right) \left(\frac{4x}{5} + \frac{3y}{4} \right)$

Using identity $(x + a)(x + b) = x^2 + (a + b)x + ab$

$$\left(\frac{4x}{5} + \frac{y}{4} \right) \left(\frac{4x}{5} + \frac{3y}{4} \right) = \left(\frac{4x}{5} \right)^2 + \left(\frac{y}{4} + \frac{3y}{4} \right) \frac{4x}{5} + \left(\frac{y}{4} \right) \times \left(\frac{3y}{4} \right)$$

$$= \frac{16}{25} x^2 + \frac{4}{5} xy + \frac{3}{16} y^2$$

ii. $(x^2 + y^2)(x^2 - y^2)$

Using identity $(a + b)(a - b) = a^2 - b^2$

$$(x^2 + y^2) = (x^2)^2 - (y^2)^2 = x^4 - y^4$$

iii. $(0.9p - 0.5q)^2$

Using identity $(a - b)^2 = a^2 - 2ab + b^2$

$$\begin{aligned} (0.9p - 0.5q)^2 &= (0.9p)^2 - 2(0.9)(0.5q) + (0.5q)^2 \\ &= 0.81p^2 - 0.9pq + 0.25q^2 \end{aligned}$$

5. Evaluate $(-7x + y) X (3x^2 + xyz + y^2)$, when $x = 1, y = 2, z = 3$.

Sol. We have $(-7x + y) X (3x^2 + xyz + y^2)$

$$\begin{aligned} &= -7x X (3x^2 + xyz + y^2) + y X (3x^2 + xyz + y^2) \\ &= -7x X 3x^2 - 7x X xyz - 7x X y^2 + y X 3x^2 + y X xyz + y X y^2 \\ &= -21x^3 - 7x^2yz - 7xy^2 + 3x^2y + xy^2z + y^3 \end{aligned}$$

Putting the value of $x = 1, y = 2, z = 3$ in the product so obtained, we get

Required value

$$\begin{aligned} &= -21 \times 1^3 - 7 \times 1^2 \times 2 \times 3 - 7 \times 1 \times 2^2 + 3 \times 1^2 \times 2 + 1 \times 2^2 \times 3 + 2^3 \\ &= -21 - 42 - 28 + 6 + 12 + 8 = -65 \end{aligned}$$

I. Long Answer Questions

1. If $a + b = 25$ and $a^2 + b^2 = 225$, then find ab .

[NCERT Exemplar]

Sol. We know that,

$$(a + b)^2 = a^2 + b^2 + 2ab$$

Here, $a + b = 25, a^2 + b^2 = 225$

$$\Rightarrow (25)^2 = 225 + 2ab$$

$$\Rightarrow 625 = 225 + 2ab$$

$$625 - 225 = 2ab$$

$$\Rightarrow 400 = 2ab$$

$$\Rightarrow ab = \frac{400}{2}$$

$$\Rightarrow ab = 200$$

2. Find the value of

$$\text{a. } \frac{6.25 \times 6.25 - 1.75 \times 1.75}{4.5}$$

$$\text{b. } \frac{198 \times 198 - 102 \times 102}{96}$$

Sol. a. $\frac{6.25 \times 6.25 - 1.75 \times 1.75}{4.5}$

$$= \frac{(6.25)^2 - (1.75)^2}{4.5}$$

$$= \frac{(6.25+1.75)(6.25-1.75)}{4.5}$$

$$[\because a^2 - b^2 = (a+b)(a-b)]$$

$$= \frac{8 \times 4.5}{4.5}$$

$$= 8$$

b. $\frac{198 \times 198 - 102 \times 102}{96}$

$$= \frac{(198)^2 - (102)^2}{96}$$

$$= \frac{(198+102)(198-102)}{96}$$

$$[\because a^2 - b^2 = (a+b)(a-b)]$$

$$= \frac{300 \times 96}{96}$$

$$= 300$$

3. Show that : $(4pq + 3q)^2 - (4pq - 3q)^2 = 48pq^2$

[NCERT Exemplar]

Sol.

$$\text{L.H.S} = (4pq + 3q)^2 - (4pq - 3q)^2$$

$$= (4pq)^2 + 2(4pq)(3q) + (3q)^2 - [(4pq)^2 - 2(4pq)(3q) + (3q)^2]$$

$$= 16p^2q^2 + 24pq^2 + 9q^2 - [16p^2q^2 - 24pq^2 + 9q^2]$$

$$= 48pq^2 = \text{R.H.S}$$

[NCERT Exemplar]

4. Subtract : $4p^2q + 5pq^2 - 3pq + 7q - 8p - 10$ from $5p^2q - 2pq^2 + 5pq - 11q - 3p + 28$.

Sol. We have,

$$5p^2q - 2pq^2 + 5pq - 11q - 3p + 28$$

$$4p^2q + 5pq^2 - 3pq + 7q - 8p - 10$$

$$\begin{array}{r} (-) \quad (-) \quad (+) \quad (-) \quad (+) \quad (+) \\ \hline p^2 - 7pq^2 + 8pq - 18q + 5p + 38 \end{array}$$

5. The length and breadth of a rectangle are $3x^2 - 2$ and $2x + 5$ respectively, then find its area.

Sol. Here, Length = $3x^2 - 2$

Breadth = $2x + 5$

Area = Length x Breadth

$$= (3x^2 - 2) \times (2x + 5)$$

$$= 3x^2(2x + 5) + (-2)(2x + 5)$$

$$= (3x^2 \times 2x) + (5 \times 3x^2) + (-2) \times 2x + (-2) \times 5$$

$$= 6x^3 + 15x^2 + (-4x) + (-10)$$

$$= 6x^3 + 15x^2 - 4x - 10$$

6. Find the volume of cuboid whose dimensions are $(x^2 - 2)$, $(2x + 2)$ and $(x - 1)$

Sol. Volume of a cuboid = Length x Breadth x Height

$$= (x^2 - 2) [(2x + 2)(x - 1)]$$

$$= (x^2 - 2) [2x^2 - 2x + 2x - 2]$$

$$= (x^2 - 2) [2x^2 - 2]$$

$$= 2(x^2 - 2)(x^2 - 1)$$

$$= 2[x^4 - x^2 - 2x^2 + 2]$$

$$= 2[x^4 - 3x^2 + 2]$$

$$= 2x^4 - 6x^2 + 4$$

II. Long Answer Questions

1. Subtract

i. $2ab^2c^3 + 4a^2b^2 c - 5a^2bc^2$ from $- 10a^2b^2 c + 4ab^2c^2 + 2a^2bc^2$

ii. $-3p^2 + 3pq + 3px$ from $3p(-p - a - r)$

Sol. (i)
$$\begin{array}{r} -10a^2b^2c & + 4ab^2c^2 & + 2a^2bc^2 \\ + 4a^2b^2c & = 2ab^2c^2 & - 5a^2bc^2 \\ (-1) & (-) & (+) \\ \hline -14a^2b^2c & + 2ab^2c^2 & + 7a^2bc^2 \end{array}$$

ii. $3p(-p - a - r) = [3p \times (-p)] + [3p \times (-1)] + [3p \times (-r)]$

$$= -3p^2 - 3pa - 3pr$$

Subtracting

$$\begin{array}{r} -3p^2 - 3pa - 3pr \\ -3p^2 & + 3pq + 3p \\ (+) & (-1) & (2) \\ \hline -3pa - 3pr - 3pq - 3px \end{array}$$

$$\text{So, } 3p(-p - a - r) - (-3p^2 + 3pq + 3px) = -3pa - 3pr - 3pq - 3px$$

2. If $x + y = 12$ and $xy = 14$, find the value of $x^2 + y^2$.

Sol. We know that $(a + b)^2 = a^2 + 2ab + b^2$

$$(x + y)^2 = x^2 + 2xy + y^2$$

Putting the values of $x + y = 12$ and $xy = 14$, we have

$$(12)^2 = x^2 + y^2 + 2 \times 14$$

$$\text{or, } 144 = x^2 + y^2 + 28$$

$$\text{or, } 144 - 28 = x^2 + y^2$$

$$\text{or, } x^2 + y^2 = 116$$

Which is the required answer.

3. If $(x + \frac{1}{x}) = 7$, find the values of $(x^2 + \frac{1}{x^2})$ and $(x^4 + \frac{1}{x^4})$.

Sol. We have, $(x + \frac{1}{x}) = 7$

$$\begin{aligned} (x + \frac{1}{x})^2 &= (7)^2 \\ \Rightarrow x^2 + \frac{1}{x^2} + 2 \times x \times \frac{1}{x} &= 49 \quad (\text{Using } (a+b)^2 = a^2 + b^2 + 2ab \text{ to the LHS}) \\ \Rightarrow x^2 + \frac{1}{x^2} + 2 &= 49 \Rightarrow x^2 + \frac{1}{x^2} = 49 - 2 \\ \Rightarrow x^2 + \frac{1}{x^2} &= 47 \quad \dots \dots (1) \end{aligned}$$

For finding the second quantity, squaring both sides in (1),

$$\begin{aligned} (x^2 + \frac{1}{x^2})^2 &= (47)^2 \\ \Rightarrow (x^2)^2 + (\frac{1}{x^2})^2 + 2 \cdot x^2 \cdot (\frac{1}{x^2}) &= 2209 \quad [\text{Using again } (a+b)^2 = a^2 + b^2 + 2ab] \\ \Rightarrow x^4 + \frac{1}{x^4} + 2 &= 2209 \Rightarrow x^4 + \frac{1}{x^4} = 2209 - 2 \\ x^4 + \frac{1}{x^4} &= 2207 \end{aligned}$$

4. If $x + \frac{1}{x} = 6$, find the value of $x^3 + \frac{1}{x^3}$

Sol. We have, $x + \frac{1}{x} = 6$

On cubing both the sides, we get $(x + \frac{1}{x})^3 = 6^3$

$$\begin{aligned} \Rightarrow x^3 + \frac{1}{x^3} + 3 \times x \times \frac{1}{x} (x + \frac{1}{x}) &= 216 \\ \Rightarrow x^3 + \frac{1}{x^3} + 3 (x + \frac{1}{x}) &= 216 \quad \dots \dots \text{(ii)} \end{aligned}$$

Substituting $x + \frac{1}{x} = 6$ [from (i)] in (ii)

$$\begin{aligned} \Rightarrow x^3 + \frac{1}{x^3} + 3 \times 6 &= 216 \Rightarrow x^3 + \frac{1}{x^3} + 18 &= 216 \\ \Rightarrow x^3 + \frac{1}{x^3} &= 216 - 18 = 198 \end{aligned}$$

5. Obtain the product of following.

$$(xy^2 - 3), \left(\frac{1}{3}x^2y^2 + 3xy + x\right)$$

Sol. We have $(xy^2 - 3), \left(\frac{1}{3}x^2y^2 + 3xy + x\right)$

Since one coefficient is in fraction, therefore row method only preferred.

$$\text{Required product} = (xy^2 - 3) \times \left(\frac{1}{3}x^2y^2 + 3xy + x\right)$$

$$\begin{aligned}
 &= xy^2 \left(\frac{1}{3}x^2 y^2 + 3xy + x \right) + (-3) \left(\frac{1}{3}x^2 y^2 + 3xy + x \right) \\
 &= xy^2 \times \frac{1}{3}x^2 y^2 + xy^2 \times 3xy + xy^2 \times x + (-3) \times \frac{1}{3}x^2 y^2 + (-3) \times 3xy + (-3)x \\
 &= \frac{1}{3}x^{1+2} y^{2+2} + 3x^{1+1} y^{2+1} + x^{1+1} y^2 - x^2 y^2 - 9xy - 3x \\
 &= \frac{1}{3}x^3 y^4 + 3x^2 y^3 + x^2 y^2 - x^2 y^2 - 9xy - 3x \\
 &= \frac{1}{3}x^3 y^4 + 3x^2 y^3 - 9xy - 3x
 \end{aligned}$$

I. Higher Order Thinking Skill

1. If $x + \frac{1}{x} = 5$, find the value of $x^2 + \frac{1}{x^2}$.

[NCERT Exemplar]

Sol. Since, $x + \frac{1}{x} = 5$

Squaring on both sides,

$$\left(x + \frac{1}{x} \right)^2 = 5^2$$

$$\text{or } x^2 + \frac{1}{x^2} + 2 \times x \times \frac{1}{x} = (5)^2$$

$$\text{or } x^2 + \frac{1}{x^2} + 2 = 25$$

$$\text{or } x^2 + \frac{1}{x^2} = 25 - 2$$

$$\text{or } x^2 + \frac{1}{x^2} = 23$$

2. Using division show that $(x + 1)$ is a factor of $(2x^2 + 3x + 1)$.

[NCERT Exemplar]

Sol. $x + 1 \overline{)2x^2 + 3x + 1}$ $(x^2 + 1)$

$$\begin{array}{r}
 2x^2 + 3x + 1 \\
 - - - \\
 x + 1 \\
 - - - \\
 x + 1 \\
 - - - \\
 0
 \end{array}$$

Since, remainder = 0

Then, $(x + 1)$ is a factor of $(2x^2 + 3x + 1)$.

I. Value Based Questions

1. a. Find the value of the expression $(81x^2 + 16y^2 - 72xy)$, when $x = \frac{2}{3}$ and $y = \frac{3}{4}$.

b. If $a = 2$ and $b = 5$, then verify $(a + b)^2 = a^2 + b^2 + 2ab$. [NCERT Exemplar]

Sol. a. $81x^2 + 16y^2 - 72xy = (9x)^2 + (4y)^2 - 2 \times 9x \times 4y$

$$= (9x - 4y)^2$$

$$[\because a^2 + b^2 - 2ab = (a - b)^2]$$

Now, putting $x = \frac{2}{3}$ and $y = \frac{3}{4}$, then

$$= \left(9 \times \frac{2}{3} - 4 \times \frac{3}{4}\right)^2$$

$$= (6 - 3)^2 = 3^2 = 9$$

b. Putting $a = 2$ and $b = 5$, then

$$\text{L.H.S} = (a + b)^2$$

$$= (2 + 5)^2 = 7^2 = 49$$

and

$$\text{R.H.S} = a^2 + b^2 + 2ab$$

$$= 2^2 + 5^2 + 2 \times 2 \times 5$$

$$= 4 + 25 + 20 = 49$$

Hence, L.H.S = R.H.S = 49



Next Generation School